arXiv:1209.5030v1 [math.NT] 23 Sep 2012

A SIMPLE COMPUTATION OF $\zeta(2k)$ BY USING BERNOULLI POLYNOMIALS AND A TELESCOPING SERIES

ÓSCAR CIAURRI, LUIS M. NAVAS, FRANCISCO J. RUIZ, AND JUAN L. VARONA

Abstract. We present a new proof of Euler's formulas for $\zeta(2k)$, where $k = 1, 2, 3, \dots$, which uses only the defining properties of the Bernoulli polynomials, obtaining the value of $\zeta(2k)$ by summing a telescoping series. Only basic techniques from Calculus are needed to carry out the computation. The method also applies to $\zeta(2k+1)$ and the harmonic numbers, yielding integral formulas for these.

1. Introduction

In the mathematical literature, one finds many ways of obtaining the formula

$$\zeta(2k) := \sum_{n=1}^{\infty} \frac{1}{n^{2k}} = \frac{(-1)^{k-1} 2^{2k-1} \pi^{2k}}{(2k)!} B_{2k}, \qquad k = 1, 2, 3, \dots, \tag{1}$$

where B_k is the kth Bernoulli number, a result first published by Euler in 1734. For example, the recent paper [1] contains quite a complete list of references. The aim of this paper is to give a new proof of (1) which is simple and elementary, in the sense that it involves only the Bernoulli polynomials and a telescoping series.

2. The Bernoulli polynomials

For completeness, we begin by recalling the definition of the Bernoulli polynomials $B_k(t)$ and their basic properties. There are of course multiple approaches one can take (see [4], which shows seven ways of defining these polynomials). A frequent starting point is the generating function

$$\frac{x e^{xt}}{e^x - 1} = \sum_{k=0}^{\infty} B_k(t) \frac{x^k}{k!}, \tag{2}$$

from which, by the uniqueness theorem for power series expansions, one can quickly obtain many of the basic properties, among which we single out

$$B_0(t) = 1, \quad B_k'(t) = k B_{k-1}(t), \qquad k \geq 1, \tag{3}$$

which by induction shows that $B_k(t)$ is in fact a polynomial, and

$$\int_0^1 B_k(t)\, dt = 0, \qquad k \geq 1. \tag{4}$$

If one considers that the generating function approach is not completely "elementary," in the sense of requiring more than standard first year Calculus (series of

---

2000 *Mathematics Subject Classification.* Primary 40C15; Secondary 11M06.

Key words and phrases. Riemann zeta function, Bernoulli polynomials, Bernoulli numbers, telescoping series.

The research of the authors is supported by grant MTM2012-36732-C03-02 of the DGI.

functions, justifying exchanging the derivative with the infinite sum, etc.) it is easily avoided by instead using (3) and (4) to define the polynomials $B_k(t)$. In any case, one finds that the first few Bernoulli polynomials are

$$B_0(t) = 1, \qquad B_1(t) = t - \frac{1}{2}, \qquad B_2(t) = t^2 - t + \frac{1}{6}.$$

The Bernoulli *numbers* are defined to be the values $B_k = B_k(0)$. From (3), (4), and the Fundamental Theorem of Calculus, one sees that $B_k(0) = B_k(1)$ for $k \geq 2$, and from the symmetry relation

$$B_k(1-t) = (-1)^k B_k(t), \qquad k \geq 0 \tag{5}$$

(easily proved by induction on k) one deduces

$$\begin{aligned} B_{2k}(1) &= B_{2k}(0) = B_{2k}, \qquad k \geq 0, \\ B_{2k+1}(1) &= B_{2k+1}(0) = 0, \qquad k \geq 1. \end{aligned} \tag{6}$$

There are many other properties and relations satisfied by the Bernoulli polynomials, but those listed above are the only ones necessary for the goal we have in mind.

3. Computation of $\zeta(2k)$

3.1. **Some auxiliary integrals.**

Consider the integrals

$$I(k,m) := \int_0^1 B_{2k}(t)\cos(m\pi t)\,dt, \qquad k \geq 0,\ m \geq 1.$$

An immediate computation shows that $I(0,m) = 0$ for $m \geq 1$, and, for $k \geq 1$, integrating by parts twice and applying (3), we get

$$\begin{aligned} I(k,m) &= \frac{1}{m\pi}\Big[B_{2k}(t)\sin(m\pi t)\Big]_{t=0}^{t=1} - \frac{2k}{m\pi}\int_0^1 B_{2k-1}(t)\sin(m\pi t)\,dt \\ &= \frac{2k}{m^2\pi^2}\Big[B_{2k-1}(t)\cos(m\pi t)\Big]_{t=0}^{t=1} - \frac{2k(2k-1)}{m^2\pi^2}\, I(k-1,m), \end{aligned}$$

which gives us both the special case

$$I(1,m) = \int_0^1 \left(t^2 - t + \frac{1}{6}\right)\cos(m\pi t)\,dt = \begin{cases} 0, & m = 1,3,5,\dots, \\ \dfrac{2}{m^2\pi^2}, & m = 2,4,6,\dots \end{cases}$$

and, by (6), the recurrence relation

$$I(k,m) = -\frac{2k(2k-1)}{m^2\pi^2}\, I(k-1,m), \qquad k \geq 2,$$

from which, by induction, one obtains the closed form

$$I(k,m) = \begin{cases} 0, & m = 1,3,5,\dots, \\ \dfrac{(-1)^{k-1}(2k)!}{m^{2k}\pi^{2k}}, & m = 2,4,6,\dots. \end{cases} \tag{7}$$

Now, for reasons which are made clear below, consider $B_k^*(t) = B_k(t) - B_k(0) = B_k(t) - B_k$, i.e., the Bernoulli polynomial minus its constant term. The corresponding integral

$$I^*(k,m) := \int_0^1 B_{2k}^*(t)\cos(m\pi t)\,dt = \int_0^1 (B_{2k}(t) - B_{2k})\cos(m\pi t)\,dt$$

is equal to $I(k,m)$, because $\int_0^1 \cos(m\pi t)\,dt = 0$. For fixed $k \geq 1$, summing $I^*(k,m)$ over $m \geq 1$ yields, by (7),

$$\begin{aligned}\frac{(-1)^{k-1}(2k)!}{2^{2k}\pi^{2k}}\zeta(2k) &= \frac{(-1)^{k-1}(2k)!}{\pi^{2k}}\sum_{m=1}^{\infty}\frac{1}{(2m)^{2k}} = \sum_{m=1}^{\infty} I^*(k,2m)\\ &= \sum_{m=1}^{\infty} I^*(k,2m) + \sum_{m=0}^{\infty} I^*(k,2m+1) = \sum_{m=1}^{\infty} I^*(k,m)\\ &= \sum_{m=1}^{\infty}\int_0^1 B_{2k}^*(t)\cos(m\pi t)\,dt.\end{aligned}$$

3.2. **The telescoping trick.** We need the elementary trigonometric identity

$$\cos(mx) = \frac{\sin(\frac{2m+1}{2}x) - \sin(\frac{2m-1}{2}x)}{2\sin(\frac{x}{2})}, \tag{8}$$

which can be proved, for example, by induction. With the introduction of (8), we now have a telescoping series, yielding

$$\begin{aligned}&\frac{(-1)^{k-1}(2k)!}{2^{2k}\pi^{2k}}\zeta(2k)\\ &\quad= \lim_{N\to\infty}\sum_{m=1}^{N}\left(\int_0^1 B_{2k}^*(t)\frac{\sin(\frac{2m+1}{2}\pi t)}{2\sin(\frac{\pi t}{2})}\,dt - \int_0^1 B_{2k}^*(t)\frac{\sin(\frac{2m-1}{2}\pi t)}{2\sin(\frac{\pi t}{2})}\,dt\right)\\ &\quad= \left(\lim_{N\to\infty}\int_0^1 B_{2k}^*(t)\frac{\sin(\frac{2N+1}{2}\pi t)}{2\sin(\frac{\pi t}{2})}\,dt\right) - \frac{1}{2}\int_0^1 B_{2k}^*(t)\,dt.\end{aligned}$$

We observe that by (4), the value of the second term is

$$\frac{1}{2}\int_0^1 B_{2k}^*(t)\,dt = \frac{1}{2}\int_0^1 (B_{2k}(t) - B_{2k})\,dt = -\frac{B_{2k}}{2}.$$

Now, we show that the limit in the first term is zero. Note that the function

$$f(t) = \frac{B_{2k}^*(t)}{2\sin(\frac{\pi t}{2})}, \qquad t \in (0,1],$$

extends by continuity to $t = 0$ since $B_{2k}^*(0) = 0$ (this is the reason for subtracting the constant term), and is differentiable on $[0,1]$ with a continuous derivative. Denoting $(2N+1)\pi/2$ by R, integrating by parts gives

$$\int_0^1 f(t)\sin(Rt)\,dt = -\frac{\cos(R)}{R}f(1) + \frac{1}{R}f(0) + \int_0^1 f'(t)\frac{\cos(Rt)}{R}\,dt.$$

The boundedness of $f'(t)$ and the trivial estimate $|\cos t| \leq 1$ show that each term in the above sum approaches zero as $R \to \infty$, so that indeed the limit tends to 0. Consequently,

$$\frac{(-1)^{k-1}(2k)!}{2^{2k}\pi^{2k}}\,\zeta(2k) = \frac{B_{2k}}{2}$$

which after rearranging terms, gives (1).

4. Fourier Confidential

Readers familiar with the basic theory of Fourier series will of course immediately recognize that the integrals $I(k,m)$ are essentially the Fourier coefficients of the Bernoulli polynomial $B_{2k}(t)$. Namely, the Fourier series of $B_{2k}(t)$, first computed by Hurwitz, is

$$B_{2k}(t) = \frac{(-1)^{k-1}(2k)!}{2^{2k-1}\pi^{2k}} \sum_{m=1}^{\infty} \frac{\cos(2\pi m x)}{m^{2k}}, \qquad t \in [0,1),\ k \geq 1.$$

and setting $t = 0$ yields (1). Of course, this approach necessitates proving the basic facts about Fourier series, not the least of which are the issue of pointwise convergence and the Riemann-Lebesgue Lemma. Fortunately, the latter is easily proved for C^1 functions with the argument using integration by parts we have given, and the issue of convergence is dealt with using the "telescoping trick" (8).

The use of the same "telescope" (8) in [2] brought to our attention that this idea, used there to compute $\zeta(2)$ via the integral $\int_0^\pi x\cos(mx)\,dx$, could also be used to compute $\zeta(2k)$. However, one needs to realize that instead of integrating $x^k\cos(mx)$, the natural functions to integrate against the cosine are the Bernoulli polynomials.

The difference formula (8) will be recognized as that obtained from the definition of the Dirichlet kernel and its expression as

$$D_N(t) = \sum_{m=-N}^{N} e^{im\pi t} = 1 + 2\sum_{m=1}^{N} \cos(m\pi t) = \frac{\sin\left(\frac{2N+1}{2}\pi t\right)}{\sin\left(\frac{\pi t}{2}\right)}.$$

It is also worthwhile noting that a related telescoping trick also yields a nice result regarding pointwise convergence of Fourier series in general, as shown in [3].

One also needs to see how the basic defining relations such as (3) can be employed to give an elementary method of computation for the Fourier coefficients. We have used the recurrence relation (3) to derive the formulas for the coefficients in a straightforward manner. This approach is actually not standard, although it seems to us to be the simplest. Most classical approaches involve the use of complex analytic summation techniques, due in part to the fact that they generalize, for example, to the Hurwitz zeta function. Such techniques are quite useful in analytic number theory, but would negate the elementary nature of the proof given here.

5. What about $\zeta(2k+1)$?

The same approach will yield a formula for $\zeta(2k+1)$, but the term which is subtracted when summing the telescoping series is an integral which, as far as anyone knows, cannot be evaluated in (a non-tautological) closed form. We proceed in the same way as before, except this time, we consider the integrals

$$J(k,m) := \int_0^1 B_{2k+1}(t)\sin(m\pi t)\,dt$$

(which are the Fourier sine coefficients). Direct computation shows that

$$J(0,m) = \int_0^1 \left(t - \frac{1}{2}\right)\sin(m\pi t)\,dt = -\frac{1+(-1)^m}{m\pi} = \begin{cases} 0, & m = 1,3,5,\ldots, \\ -\dfrac{1}{m\pi}, & m = 2,4,6,\ldots \end{cases}$$

and, for $k \geq 1$, integrating by parts twice gives, using (3) and (6), the recurrence relation

$$J(k,m) = -\frac{(2k+1)(2k)}{m^2\pi^2}\, J(k-1,m)$$

from which, by induction, we obtain the closed form

$$J(k,m) = \begin{cases} 0, & m = 1,3,5,\ldots, \\ \dfrac{(-1)^{k-1}(2k+1)!}{m^{2k+1}\pi^{2k+1}}, & m = 2,4,6,\ldots. \end{cases} \tag{9}$$

Note that this time, since $B_{2k+1}(0) = B_{2k+1} = 0$ for $k \geq 1$, subtracting the constant term is not necessary except for $k = 0$, since $B_1 = -1/2$, although, since $\zeta(1) = \infty$, this is irrelevant for now (but see the next section). Thus, summing (9) over m, we obtain, for $k \geq 1$,

$$\begin{aligned} \frac{(-1)^{k-1}(2k+1)!}{2^{2k+1}\pi^{2k+1}}\,\zeta(2k+1) &= \frac{(-1)^{k-1}(2k+1)!}{\pi^{2k+1}} \sum_{m=1}^{\infty} \frac{1}{(2m)^{2k+1}} = \sum_{m=1}^{\infty} J(k,2m) \\ &= \sum_{m=1}^{\infty} J(k,2m) + \sum_{m=0}^{\infty} J(k,2m+1) = \sum_{m=1}^{\infty} J(k,m) \\ &= \sum_{m=1}^{\infty} \int_0^1 B_{2k+1}(t)\sin(m\pi t)\,dt. \end{aligned}$$

The trigonometric identity that plays the role of (8) in this case is

$$\sin(mx) = -\frac{\cos(\frac{2m+1}{2}x) - \cos(\frac{2m-1}{2}x)}{2\sin(\frac{x}{2})}. \tag{10}$$

Thus, we now obtain the telescoping series

$$\begin{aligned} &\frac{(-1)^{k-1}(2k+1)!}{2^{2k+1}\pi^{2k+1}}\,\zeta(2k+1) \\ &= -\lim_{N\to\infty} \sum_{m=1}^{N} \left(\int_0^1 B_{2k+1}(t)\frac{\cos(\frac{2m+1}{2}\pi t)}{2\sin(\frac{\pi t}{2})}\,dt - \int_0^1 B_{2k+1}(t)\frac{\cos(\frac{2m-1}{2}\pi t)}{2\sin(\frac{\pi t}{2})}\,dt \right) \\ &= -\left(\lim_{N\to\infty} \int_0^1 B_{2k+1}(t)\frac{\cos(\frac{2N+1}{2}\pi t)}{2\sin(\frac{\pi t}{2})}\,dt \right) + \int_0^1 B_{2k+1}(t)\frac{\cos(\frac{\pi t}{2})}{2\sin(\frac{\pi t}{2})}\,dt. \end{aligned}$$

The limit is null, for the same reason as before. Consequently, we have proved that

$$\zeta(2k+1) = \frac{(-1)^{k-1}2^{2k}\pi^{2k+1}}{(2k+1)!} \int_0^1 B_{2k+1}(t)\cot\left(\frac{\pi t}{2}\right)dt, \qquad k \geq 1. \tag{11}$$

For example,

$$\zeta(3) = \frac{\pi^3}{3} \int_0^1 \left(2t^3 - 3t^2 + t\right)\cot\left(\frac{\pi t}{2}\right)dt.$$

It would be nice to know if the integral in (11) has a closed form expression (other than in terms of $\zeta(2k+1)$ of course!) but at present this problem remains open.

6. $\zeta(1)$: THE HARMONIC SERIES

Although the harmonic series diverges, we can still obtain information from the integrals $J(0,m)$ if we consider the partial sums of the telescoping series in the last section with $k=0$. It is actually easier *not* to remove the constant term $B_1=-1/2$, but if we don't, note that each difference of integrals in the telescoping sum is now an indeterminacy of the form $\infty-\infty$. The trick is to just use (10) to sum $\sin(m\pi x)$ *inside* the integral $J(0,m)$. Then it is easy to see that we obtain the formula

$$\int_0^1 \left(t-\frac{1}{2}\right)\frac{\cos\left(\frac{4M+1}{2}\pi t\right)-\cos\left(\frac{\pi t}{2}\right)}{\sin\left(\frac{\pi t}{2}\right)}\,dt = \frac{1}{\pi}\,H_M, \qquad M\ge 1, \tag{12}$$

where $H_M=\sum_{m=1}^{M}\frac{1}{m}$ is the Mth harmonic number.

By (11), the same formula is valid with $4M+3$ instead of $4M+1$, but for $4M$ and $4M+2$, the interested reader may verify that we get combinations involving π, $\log 2$, and Catalan's constant

$$G=\sum_{n=0}^{\infty}\frac{(-1)^n}{(2n+1)^2},$$

via the formulas

$$\int_0^1 \frac{t}{\sin\left(\frac{\pi t}{2}\right)}\,dt=\frac{8}{\pi^2}\,G, \qquad \int_0^1 t\cot\left(\frac{\pi t}{2}\right)dt=\frac{2\log 2}{\pi},$$

but these, though certainly interesting, are not really part of this story.

REFERENCES

[1] E. de Amo, M. Díaz Carrillo, and J. Fernández-Sánchez, Another proof of Euler's formula for $\zeta(2k)$, *Proc. Amer. Math. Soc.* **139** (2011), no. 4, 1441–1444.

[2] D. Benko, The Basel problem as a telescoping series, *College Math. J.* **43** (2012), no. 3, 244–250.

[3] P. Chernoff, Pointwise Convergence of Fourier Series, *Amer. Math. Monthly* **87**, no. 5, 399–400.

[4] F. Costabile, F. Dell'Accio, and M. I. Gualtieri, A new approach to Bernoulli polynomials, *Rend. Mat. Appl. (7)* **26** (2006), 1–12.

Dpto. de Matemáticas y Computación, Universidad de La Rioja, 26004 Logroño, Spain
E-mail address: `oscar.ciaurri@unirioja.es`

Departamento de Matemáticas, Universidad de Salamanca, 37008 Salamanca, Spain
E-mail address: `navas@usal.es`

Dpto. de Matemáticas, Universidad de Zaragoza, 50009 Zaragoza, Spain
E-mail address: `fjruiz@unizar.es`

Dpto. de Matemáticas y Computación, Universidad de La Rioja, 26004 Logroño, Spain
E-mail address: `jvarona@unirioja.es`
URL: `http://www.unirioja.es/cu/jvarona/welcome.html`